

Profile of glycated-hemoglobin, antioxidant vitamin and cytokine levels in pulmonary tuberculosis patients: A cross sectional study at Pulmonary Diseases Center Semarang City, Indonesia

by Lintang Saraswati

Submission date: 28-Sep-2018 05:12PM (UTC+0700)

Submission ID: 1010004070

File name: Lampiran_C1-2_Artikel.pdf (278.8K)

Word count: 4879

Character count: 27279

Available online at www.sciencedirect.com

ScienceDirect

Biomedical Journal

journal homepage: www.elsevier.com/locate/bj

Original Article

Profile of glycated-hemoglobin, antioxidant vitamin and cytokine levels in pulmonary tuberculosis patients: A cross sectional study at Pulmonary Diseases Center Semarang City, Indonesia

Praba Ginandjar ^{a,*}, Lintang Dian Saraswati ^a, Bagoes Widjanarko ^{b,c}^a Department of Epidemiology and Tropical Diseases, Faculty of Public Health, Diponegoro University, Indonesia^b Department of Health Promotion, Faculty of Public Health, Diponegoro University, Indonesia^c Tuberculosis Operational Research Group (TORG), Indonesia

ARTICLE INFO

Article history:

Received 29 June 2015

Accepted 21 January 2016

Available online 9 November 2016

Keywords:

HbA1c

Pulmonary tuberculosis

Vitamin A

Vitamin E

ABSTRACT

Background: Uncontrolled blood glucose, which marked by high level of HbA1c, increases risk of pulmonary TB because of cellular immunity dysfunction. This study aimed to analyze profile of glycated hemoglobin, antioxidant vitamins status and cytokines levels in active pulmonary TB patients.

Methods: This was a cross sectional study, conducted at Pulmonary Diseases Center Semarang City, Indonesia. Study subject consisted of 62 pulmonary TB patients, diagnosed with positive acid fast bacilli and chest X-ray. ELISA was used to measure IFN- γ and IL-12. Status of antioxidant vitamins was determined by concentration of vitamin A and E using HPLC. Blood glucose control was determined by HbA1c concentration (HbA1c $\geq 7\%$ is considered as uncontrolled).

Results: A significant difference of age between pulmonary tuberculosis patients with normal and uncontrolled blood glucose ($p = 0.000$) was showed, while all other characteristics (sex, education, occupation) did not differ with $p = 0.050, 0.280, 0.380$ respectively. Mean HbA1c was $7.25 \pm 2.70\%$. Prevalence of uncontrolled glucose among pulmonary TB patients was 29%. Levels of IFN- γ and IL-12 did not differ according to HbA1c concentration ($p = 0.159$ and $p = 0.965$ respectively). Pulmonary tuberculosis patients with uncontrolled blood glucose has higher vitamin E ($p = 0.006$), while vitamin A did not differ significantly ($p = 0.478$).

Conclusions: This study supports the importance of performing diabetes screening among pulmonary TB patients. Further study needs to be done to determine the feasibility of TB-DM co-management.

* Corresponding author. Department of Epidemiology and Tropical Diseases, Faculty of Public Health, Diponegoro University, Jl. Prof. Soedarto SH, Kampus UNDIP Tembalang, Semarang 50275, Indonesia. Tel.: +62 81325887942; fax: +62 247460044ext.102.

E-mail address: praba.ginandjar@live.undip.ac.id (P. Ginandjar).

Peer review under responsibility of Chang Gung University.

<http://dx.doi.org/10.1016/j.bj.2016.01.011>

2319-4170/© 2016 Chang Gung University. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

At a glance commentary

Scientific background on the subject

Indonesia has a second largest burden of tuberculosis globally, and had 10 million cases of diabetes in 2015. Relationship of tuberculosis and diabetes is more prominent in developing countries, where TB is endemic and the burden of diabetes mellitus is increasing. Early detection uncontrolled glucose and antioxidant vitamins in pulmonary tuberculosis patients may give benefit to control the diseases.

What this study adds to the field

Almost one third of PTB patients had HbA1c levels >7%. The group of pulmonary tuberculosis with high HbA1c levels had similar levels of vitamin A, IFN- γ , and IL-12. On the other hand, vitamin E levels was higher in HbA1c > 7. Management of co-morbidities need to prioritize at risk group. This study supports the importance of performing blood glucose screening among pulmonary tuberculosis patients.

Glycated hemoglobin, commonly known as HbA1c, reflects an average blood glucose levels over a period of weeks or months. The International Expert Committee has recommended HbA1c for diabetes diagnosis [1]. Worldwide, the International Diabetes Federation (IDF) predicts that the number of people with diabetes will rise by 55% in the next 20 years [2]. The current pandemic of type 2 diabetes mellitus is accelerating in a world where approximately one third of the population is latently infected with *Mycobacterium tuberculosis* [3]. Diabetes patients have 2–5 times higher risk to develop tuberculosis compared to non-diabetes [4,5]. Once infected, diabetes patients need longer treatment and are at risk of drug resistance [6] and treatment failure [7]. Furthermore, a study in Indonesia revealed pulmonary tuberculosis patients with co-existence of diabetes had more clinical signs compared to non-diabetes [8].

Uncontrolled blood glucose, which marked by high level of glycated hemoglobin (HbA1c) [9], is one of TB risk factors [10]. Risk of active tuberculosis development in population with diabetes [4,11,12] may relate to lower antimicrobial peptide gene expression [13]. The lower antimicrobial peptide gene expression may also increase the risk of tuberculosis reactivation [13,14] and modifies clinical sign of pulmonary tuberculosis [15]. Diabetes patients are also prone to pulmonary TB infection due to phagocytosis [16] and cellular immunity [17] dysfunctions, the two main immune responses to *M. tuberculosis* [18].

Pulmonary tuberculosis and diabetes are often associated with malnutrition. The link between nutritional status and tuberculosis has long been appreciated, but remains an emerging area of study that has focused on investigations of related biomarkers and nutrient supplementation trials [19].

Pulmonary tuberculosis patients frequently suffer from deficiencies in antioxidant micronutrients, such as vitamins A and E. Both are fundamental vitamins to the integrity of the immune response, especially the host's immune response against *M. tuberculosis*. Furthermore, vitamins play an important role in glucose metabolism [20]. Vitamins also related with active pulmonary tuberculosis. A previous study showed that the levels of vitamin C and vitamin E were significantly decreased in *M. tuberculosis* patients when compared to the healthy controls [21].

Both diabetes and pulmonary tuberculosis patients are usually underdiagnosed, or delayed diagnosed [22]. Early detection of diabetes through blood glucose indicator in pulmonary tuberculosis patients, along with the underlying factors might play role in co-management of the diseases in question. In this study, we analyzed the cytokines (IFN- γ and IL-12), antioxidant vitamin (vitamin A and E), and blood glucose control (HbA1c) in active pulmonary TB patients.

Methods

Study subject

Subject was pulmonary tuberculosis patients, who sought treatment at Pulmonary Health Center (called BKPM = Balai Kesehatan Paru Masyarakat) Semarang City, Indonesia [Fig. 1]. Patients were selected by pulmonary tuberculosis symptoms as follow: persistent cough for more than 3 weeks, produce sputum, and loss weight. Two same-day sputum specimens were collected and examined for sputum smear microscopy by Ziehl Neelsen staining, and detected to have acid fast bacilli on sputum examination [23]. Patients also underwent chest X-ray to examine the existence of tuberculosis cavities. HIV serology test was done to all pulmonary tuberculosis suspects for exclusion criteria.

Eligible subjects were selected consecutively during June 11–August 23, 2014. During the course of study, there were 85 patients diagnosed as pulmonary tuberculosis, 23 of them either refused to join the study or experienced difficulties during blood collection. Overall, there were 62 pulmonary tuberculosis patients became our study subject. All subjects agreed to join the study by signing written informed consent. Ethical clearance was obtained from the Commission of Ethics of Medical and Public Health Research, Faculty of Public Health, Diponegoro University (number: 298/EC/FKM/2013).

Data collection

Variables in this study were pulmonary tuberculosis occurrence, body mass index (BMI), IFN- γ , IL-12, vitamin A, vitamin E, HbA1c levels, characteristics (age, sex, education, occupation), and history of diseases (diabetes, high blood pressure, hypercholesterolemia). Pulmonary TB referred to patients with *M. tuberculosis* infection, which is characterized by detection of acid fast bacilli in sputum using Ziehl Neelsen examination and chest X-ray. Body mass index was derived from ratio of body weight in kilogram to body height in centimeter. Education was categorized as basic




Fig. 1 – Flow diagram of study subject participation.

(elementary to junior high school) and high education (senior high school and college). Occupation was categorized as employed and unemployed.

IFN- γ and IL-12 were examined using Quantikine HS ELISA kits and read in 450 nm. Vitamin A was measured in the form of retinol serum concentration indicator, using HPLC examination and stated in $\mu\text{g/L}$. Vitamin A below 200 $\mu\text{g/L}$ is considered as deficiency. Vitamin E was measured in the form of alpha tocopherol, which preferentially absorbed and accumulated in human. The measurement also used HPLC, which stated in mg/L , and value <3 mg/L is considered as deficiency. HbA1c levels was measured as mean of blood glucose concentration in form of glycated hemoglobin using HPLC examination in %. HbA1c indicates blood glucose control within previous three months, and is considered as diabetes if it exceeds recommendation borderline of 7%. All laboratory examination was conducted at R&E Unit, Prodia Laboratory, Indonesia.

Data analysis

Distribution of data was determined by Kolmogorov–Smirnov test. Independent t-test was used to analyze data the difference of IFN- γ and IL-12, vitamin A and vitamin E concentration in pulmonary tuberculosis patients with uncontrolled and normal blood glucose levels. The difference of characteristics was analyzed with chi-square (education, occupation, history of diseases) and independent t-test (age).

Results

Our study showed 18 out of 62 (29.0%) pulmonary tuberculosis patients had uncontrolled blood glucose, which was indicated by HbA1c levels >7% [Table 1]. There was a significant difference between age ($p = 0.000$) and diabetes history ($p = 0.000$) of pulmonary tuberculosis patients with high and normal glycated hemoglobin levels, while all other variables did not

differ significantly from each other [Table 2]. Mean HbA1c levels of all subjects was 7.25%. Mean of vitamin A from all subjects (307.29 $\mu\text{g/L}$) were considered as normal (200–800 $\mu\text{g/L}$). Mean concentration of vitamin E of all subjects (14.94 mg/L) was slightly higher than normal (8–14 mg/L) [Table 3].

Pulmonary tuberculosis patients with high HbA1c levels had similar vitamin A, IFN- γ and IL-12 with those with normal HbA1c. Statistical analysis also proved the vitamin A, IFN- γ and IL-12 did not differ significantly between both groups ($p = 0.159$, 0.965 and 0.478 respectively). On the other hand, pulmonary tuberculosis patients with HbA1c $\geq 7\%$ had higher vitamin E concentration (16.78 ± 3.49 mg/L) compared to the normal group (14.18 ± 3.17 mg/L). The difference of vitamin E levels was statistically significant ($p = 0.006$).

Discussion

Our study showed most of pulmonary tuberculosis patients were male, i.e. 67.7% (in all subjects), 77.8% (in normal HbA1c group) and 63.6% (in high HbA1c group). This result is in accordance with previous descriptive epidemiology studies in Indonesia [24] and worldwide [25–27]. Men seem to be more affected than women, with a male/female ratio of 1.9 ± 0.6 for the worldwide case notification rate [28]. The global tuberculosis epidemic is also characterized by significant differences in prevalence between men and women. Rates of tuberculosis are much higher among men than women in large areas of the world, and it has been extensively documented [29].

Table 1 – Prevalence of blood glucose control among pulmonary tuberculosis patients.

HbA1c levels	n = 62	%
$\geq 7\%$	18	29.0
<7%	44	71.0

Table 2 – Characteristics of pulmonary tuberculosis patients.

Variables	All subject (n = 62)	HbA1c $\geq 7\%$ (n = 18)	HbA1c $< 7\%$ (n = 44)	p
Age (years)	40.06 \pm 15.46	52.72 \pm 9.27	34.89 \pm 14.51	0.000 ^a
BMI (kg/m ²)	19.19 \pm 3.22	20.44 \pm 3.84	18.68 \pm 2.83	0.050 ^a
Sex: male	67.7%	63.6%	77.8%	0.280 ^b
Education: basic	58.1%	54.5%	66.7%	0.380 ^b
Occupation: unemployed	27.5%	31.8%	16.7%	0.225 ^b
History of diabetes	21.0%	66.7%	2.3%	0.000 ^b
History of high blood pressure	12.9%	11.1%	13.6%	0.788 ^b
History of hypercholesterolemia	6.5%	16.7%	2.3%	0.127 ^b

^a Independent t-test.^b Chi-square test.

Age of subjects was mostly in productive ages, with most of them were 17–35 years of old. Descriptive epidemiology studies have previously found the same pattern [30,31]. Recent study revealed polymorphism in mannose lectin (MBL) is the factor affected high case of tuberculosis in productive ages [32]. When associated with HbA1c levels, our result indicated mean age of pulmonary tuberculosis patients with HbA1c $\geq 7\%$ (52.72 \pm 9.27 years) was higher than those with normal HbA1c (34.89 \pm 14.51 years). A previous study showed that tuberculosis patients with diabetes condition tends to found in older age [33]. We suggest health institutions with tuberculosis service to pay attention on patients with higher risk of comorbidities such as diabetes. Identification of the high risk groups allows the health institutions to prioritize age groups that needs to gain more attention in management of comorbidities, in this case tuberculosis and diabetes.

Prevalence of pulmonary tuberculosis patients who have uncontrolled blood glucose with HbA1c $\geq 7\%$ was 29.0%. The prevalence is in accordance with previous report that one third of the diabetes population is latently infected with *M. tuberculosis* [3]. Our study revealed most of subject with HbA1c $\geq 7\%$ had history of diabetes (12 out of 18 subjects). This needs attention since impaired glucose tolerance is a significant risk factor for developing diabetes [34]. A previous study showed tuberculosis subjects had more comorbidities than population control subjects, including diabetes [25]. The relationship between diabetes and tuberculosis is bi-directional [12]. This study supports the importance of performing diabetes screening among pulmonary TB patients. Further study needs to be done to determine the feasibility and cost-effectiveness of co-management of TB-DM.

Our study showed a higher concentration of vitamin E in pulmonary tuberculosis patients with HbA1c $\geq 7\%$. Vitamin E is the most important lipid phase chain breaking antioxidants,

and is central to the prevention of lipid peroxidation [21]. Tuberculosis patients usually demonstrate high levels of lipid peroxidase and low concentration of plasma vitamin E [35]. Lipid peroxidation has also been considered as an early marker for diabetes [36].

Previous studies observed that vitamin E was significantly depleted in tuberculosis patients when compared with healthy individuals [21,37]. Administration of micronutrients, including vitamin E, has been shown to enhance antioxidant status [35] and accelerate tuberculosis healing [38]. Vitamin E presumably will also give benefit to tuberculosis patients with diabetes. An *in vivo* study revealed supplementation with vitamin E significantly reduced glycemia and glycated hemoglobin (HbA1c) values in diabetic rats [39]. A similar result was also shown from an experiment study in diabetic patients, which demonstrated that administration of vitamin E decreased HbA1c levels in patients with inadequate glycemic control (HbA1c $\geq 8\%$) and low vitamin E levels [40].

Our study found vitamin A did not differ between pulmonary tuberculosis patients with normal and inadequate blood glucose control. The result was in contrary with a previous study that showed tuberculosis and diabetes often coincide with micronutrient deficiency, including vitamin A [20]. Several studies worldwide observed low level of vitamin A in tuberculosis patients [41–43]. However, the effect of vitamin A supplementation on tuberculosis patients in Indonesia varies according to nutritional status of the patients. In patients with good nutritional status, supplementation of vitamin A resulted in earlier sputum smear conversion [44]. On the contrary, vitamin A supplementation failed to reduce sputum conversion time in severely malnourished patients [45].

Various studies have shown the benefits of both vitamin E and vitamin A in tuberculosis patients, or in diabetes patients separately. Yet to the best of our knowledge, there has been no

Table 3 – Status of HbA1c, cytokines and vitamins of pulmonary tuberculosis patients.

Variables	Mean \pm SD (n = 62)		Min–max	p ^a
HbA1c (%)	7.25 \pm 2.70		4.4–14.4	
		HbA1c $\geq 7\%$	HbA1c $< 7\%$	
IFN- γ	19.14 \pm 13.31	22.76 \pm 12.02	17.63 \pm 13.66	0.159
IL-12	2.58 \pm 1.63	2.57 \pm 1.57	2.58 \pm 1.67	0.965
Vitamin A (μ g/L)	307.29 \pm 149.55	328.78 \pm 184.09	298.50 \pm 134.37	0.478
Vitamin E (mg/L)	14.94 \pm 3.45	16.78 \pm 3.49	14.18 \pm 3.17	0.006

^a Independent t-test.

research to study the effect of vitamin E or vitamin A in tuberculosis with inadequate glycemic control (high HbA1c). Therefore, comparison with existing literature could not be made. We found Wang et al. (2013) conducted a randomized controlled trial to determine the effect of vitamin A in the form of retinol on treatment of active pulmonary tuberculosis [20], but the result has not been published.

Our study also did not found difference of IL-12 and IFN- γ in pulmonary tuberculosis patients with normal or HbA1c $\geq 7\%$. This is possibly due to both tuberculosis and uncontrolled glucose contribute in decreasing IL-12 and IFN- γ , thus resulting in similar cytokines profile. A longitudinal study by Tsukaguchi showed tuberculosis patients whose blood glucose was under control in six months will undergo an improvement in IFN- γ production. On the contrary, IFN- γ levels in tuberculosis patient with high or uncontrolled blood glucose remained low. This means production of IFN- γ is more related to blood glucose status than to history of diabetes [46].

T cell-mediated immune responses are important in the host control of *M. tuberculosis* infection. This is evident from the increased risk of tuberculosis in individuals with deficiencies in their IFN γ and IL-12; which promotes TH1 cell differentiation signaling pathways [47]. When associated with blood glucose, the results from previous studies were contradictory. Gomez (2013) showed monocytes from diabetic patients express the decrease of IL-12 and IFN- γ secretion *in vitro*, but the relation was affected by interaction with bacteria [48]. Chu (2015) revealed glucose increases IL-12 production in stimulated PBMCs of diabetes patients through increased IL-12 gene expression [49]. An experiment in streptozotocin-induced mouse showed that *M. tuberculosis* infection causes IL-12 and IFN- γ to decrease [18]. A metabolomic study to determine whether there is a direct functional connection between serum abundance of metabolites and of immune mediators found significantly elevated serum concentrations of IFN- γ and soluble alpha chain of the IL-2 receptor alpha in TB active patients as compared to the two healthy groups [50].

There were several limitation of this study. The diagnosis of pulmonary tuberculosis did not include culture. This was related to tuberculosis program in Indonesia that only uses fast acid bacilli detection and X-ray. In addition, history of diabetes was only determined from interview with subjects and medical record if any, not by fasting and postprandial glucose serum levels. The level of nonresponse is also a concern. Proportion of eligible subjects who refused to participate in this study was 27% (23 out of 85 eligible patients). Therefore, generalization must be taken with caution due to sample size. Besides, this study used cross sectional design, which cause difficulty to establish relation between outcome and exposure of long duration like pulmonary tuberculosis and high blood glucose. A longitudinal design should be considered to investigate the role of underlying factors of pulmonary tuberculosis in conjunction with HbA1c levels.

Conclusions

Nearly one third of pulmonary tuberculosis patients in this study (29.0%) contracted with uncontrolled blood glucose,

which was indicated by HbA1c levels $>7\%$. Overall, mean HbA1c was 7.25%. Pulmonary tuberculosis patients with HbA1c $\geq 7\%$ had higher age, BMI, IFN- γ , vitamin A and vitamin E levels. However, only vitamin E that significantly difference in pulmonary tuberculosis patients according to HbA1c levels.

Identification of high risk groups allows the health institutions to prioritize age groups that needs to gain more attention in management of co-morbidities, in this case tuberculosis and diabetes. This study supports the importance of performing diabetes screening among pulmonary TB patients. Further study needs to be done to determine the feasibility and cost-effectiveness of co-management of TB-DM.

Conflicts of interest

We declare that we have no conflict of interest.

Acknowledgement

This study was supported by the Directorate General of the Higher Education Ministry of Education and Culture and the Research Institute of Diponegoro University (grant number 180-40/UN7.3.9/PG/2014). The sponsor had no role in the study design, collection, analysis or interpretation of data.

The authors thank to Pulmonary Health Center of Semarang, Central Java, Indonesia for facilitate the study and to Mr. Muh. Fauzi for assistance with participant enrollment.

REFERENCES

- [1] The International Expert Committee. International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes Care* 2009;32:1327–34.
- [2] International Diabetes Federation. IDF diabetes atlas. 6th ed. Brussels, Belgium: IDF; 2014.
- [3] World Health Organization. Tuberculosis global facts 2011/2012. WHO Stop TB Dep. 2012. p. 1–2.
- [4] Jeon CY, Murray MB. Diabetes mellitus increases the risk of active tuberculosis: a systematic review of 13 observational studies. *PLoS Med* 2008;5:e152.
- [5] Stevenson CR, Forouhi NG, Roglic G, Williams BG, Lauer JA, Dye C, et al. Diabetes and tuberculosis: the impact of the diabetes epidemic on tuberculosis incidence. *BMC Public Health* 2007;7:234.
- [6] Chang JT, Dou HY, Yen CL, Wu YH, Huang RM, Lin HJ, et al. Effect of type 2 diabetes mellitus on the clinical severity and treatment outcome in patients with pulmonary tuberculosis: a potential role in the emergence of multidrug-resistance. *J Formos Med Assoc* 2011;110:372–81.
- [7] Dooley KE, Tang T, Golub JE, Dorman SE, Cronin W. Impact of diabetes mellitus on treatment outcomes of patients with active tuberculosis. *Am J Trop Med Hyg* 2009;80:634–9.
- [8] Alisjahbana B, Sahiratmadja E, Nelwan EJ, Purwa AM, Ahmad Y, Ottenhoff THM, et al. The effect of type 2 diabetes mellitus on the presentation and treatment response of pulmonary tuberculosis. *Clin Infect Dis* 2007;45:428–35.
- [9] Huang IC, Wang PW, Liu RT, Tung SC, Chen JF, Kuo MC, et al. The influence of self-monitoring blood glucose frequency on

- the oscillation of hemoglobin A1c and chronic complications. *Chang Gung Med J* 2012;35:46–53.
- [10] Dooley KE, Chaisson RE. Tuberculosis and diabetes mellitus: convergence of two epidemics. *Lancet Infect Dis* 2009;9:737–46.
 - [11] Jabbar A, Hussain SF, Khan AA. Clinical characteristics of pulmonary tuberculosis in adult Pakistani patients with co-existing diabetes mellitus. *East Mediterr Health J* 2006;12:522–7.
 - [12] Jeon CY, Harries AD, Baker MA, Hart JE, Kapur A, Lönnroth K, et al. Bi-directional screening for tuberculosis and diabetes: a systematic review. *Trop Med Int Health* 2010;15:1300–14.
 - [13] Gonzalez-Curiel I, Castañeda-Delgado J, Lopez-Lopez N, Araujo Z, Hernandez-Pando R, Gandara-Jasso B, et al. Differential expression of antimicrobial peptides in active and latent tuberculosis and its relationship with diabetes mellitus. *Hum Immunol* 2011;72:656–62.
 - [14] Yasar K, Pehlivanoglu F, Sengöz A, Sengoz G. Coexistence of advanced age and female gender in diabetics with extrapulmonary tuberculosis: four culture-proven cases. *Gend Med* 2011;8:334–8.
 - [15] Singla R, Khan N, Al-Sharif N, Al-Sayegh MO, Shaikh MA, Osman MM. Influence of diabetes on manifestations and treatment outcome of pulmonary TB patients. *Int J Tuberc Lung Dis* 2006;10:74–9.
 - [16] Ljubic S, Balachandran A, Pavlic-Renar I, Barada A, Metelko Z. Pulmonary infections in diabetes mellitus. *Diabetol Croat* 2005;33:115–24.
 - [17] Chang F, Shao M. Decreased cell-mediated immunity in patients with non-insulin-dependent diabetes mellitus. *Diabetes Res Clin Pract* 1995;28:137–46.
 - [18] Yamashiro S, Kawakami K, Uezu K, Kinjo T, Miyagi K, Nakamura K, et al. Lower expression of Th1-related cytokines and inducible nitric oxide synthase in mice with streptozotocin-induced diabetes mellitus infected with *Mycobacterium tuberculosis*. *Clin Exp Immunol* 2005;139:57–64.
 - [19] Frediani JK, Tukvadze N, Sanikidze E, Kipiani M, Hebbbar G, Easley KA, et al. A culture-specific nutrient intake assessment instrument in patients with pulmonary tuberculosis. *Clin Nutr* 2013;32:1023–8.
 - [20] Wang Q, Ma A, Bygbjerg JC, Han X, Liu Y, Zhao S, et al. Rationale and design of a randomized controlled trial of the effect of retinol and vitamin D supplementation on treatment in active pulmonary tuberculosis patients with diabetes. *BMC Infect Dis* 2013;13:104.
 - [21] Johnkenedy N, Onyinyechi AS, Chukwunyere NE. The antioxidant status and lipid peroxidation product of newly diagnosed and 6 weeks follow-up patients with pulmonary tuberculosis in Owerri, Imo state, Nigeria. *Asian Pac J Trop Dis* 2011;1:292–4.
 - [22] Lawn SD. Diagnosis of pulmonary tuberculosis. *Curr Opin Pulm Med* 2013;19:280–8.
 - [23] Depkes RI. Standar reagen ziehl neelsen. Jakarta: Direktorat Bina Pelayanan Penunjang Medik; 2008.
 - [24] Nasution EJS. Profil penderita tuberkulosis paru dengan diabetes mellitus dihubungkan dengan kadar gula darah puasa. Universitas Sumatera Utara; 2007.
 - [25] Legaard A, Riis A, Kornum JB, Prahl JB, Thomsen VØ, Sørensen HT, et al. Diabetes, glycemic control, and risk of tuberculosis: a population-based case control study. *Diabetes Care* 2011;34:2530–5.
 - [26] Mirsaedi MS, Tabarsi P, Radpour O, Mansouri D, Amiri M, Bagheri Z, et al. Differences in characteristics between Afghani and Iranian patients with pulmonary tuberculosis. *Int J Infect Dis* 2007;11:180–2.
 - [27] Dujaili JA, Syed Sulaiman SA, Hassali MA, Blebil AQ, Awaisu A. Socio-demographic characteristics related to quality of life in patients with pulmonary tuberculosis in Baghdad, Iraq. *Value Health* 2014;17:A280.
 - [28] World Health Organization. Global tuberculosis report 2014. Geneva, Switzerland: World Health Organization; 2014.
 - [29] Rhines AS. The role of sex differences in the prevalence and transmission of tuberculosis. *Tuberculosis (Edinb)* 2013;93:104–7.
 - [30] Ohmori M, Yoshiyama T, Ishikawa N. Estimation of the future epidemiological situation of tuberculosis in Japan. *Kekkaku* 2008;83:365–77.
 - [31] Yu M-C, Bai K-J, Chang J-H, Lee C-N. Age transition of tuberculosis patients in Taiwan, 1957–2001. *J Formos Med Assoc* 2006;105:25–30.
 - [32] Hijikata M, Matsushita I, Hang NT, Maeda S, Thuong PH, Tam DB, et al. Age-dependent association of mannose-binding lectin polymorphisms with the development of pulmonary tuberculosis in Viet Nam. *Hum Immunol* 2014;75:840–6.
 - [33] Magee MJ, Bloss E, Shin SS, Contreras C, Huaman HA, Ticona JC, et al. Clinical characteristics, drug resistance, and treatment outcomes among tuberculosis patients with diabetes in Peru. *Int J Infect Dis* 2013;17:e404–12.
 - [34] Niazi AK, Kalra S. Diabetes and tuberculosis: a review of the role of optimal glycemic control. *J Diabetes Metab Disord* 2012;11:28.
 - [35] Seyedrezadeh E, Ostadrahimi A, Mahboob S, Assadi Y, Ghaemmagami J, Pourmogaddam M. Effect of vitamin E and selenium supplementation on oxidative stress status in pulmonary tuberculosis patients. *Respirology* 2008;13:294–8.
 - [36] Arora R, Vig AP, Arora S. Lipid peroxidation: a possible marker for diabetes. *J Diabetes Metab* 2013;S11:1–6.
 - [37] Akiibinu M, Arinola O, Ogunlewe J, Onih E. Non-enzymatic antioxidants and nutritional profiles in newly diagnosed pulmonary tuberculosis patients in Nigeria. *Afr J Biomed Res* 2010;10:223–8.
 - [38] Turchenko L, Voloshchuk E, Ivanov V, Kalinovskiy T, Niedzwiecki A, Rath M. Clinical improvement of active tuberculosis patients with complex treatment and nutritional supplementation. *Open Nat Prod J* 2009;1:20–6.
 - [39] Roldi LP, Pereira RVF, Tronchini EA, Rizo GV, Scoaris CR, Zanon JN, et al. Vitamin E (alpha-tocopherol) supplementation in diabetic rats: effects on the proximal colon. *BMC Gastroenterol* 2009;9:88.
 - [40] Suksomboon N, Poolsup N, Sinprasert S. Effects of vitamin E supplementation on glycaemic control in type 2 diabetes: systematic review of randomized controlled trials. *J Clin Pharm Ther* 2011;36:53–63.
 - [41] Ramachandran G, Santha T, Garg R, Baskaran D, Iliayas SA, Venkatesan P, et al. Vitamin A levels in sputum-positive pulmonary tuberculosis patients in comparison with household contacts and healthy “normals”. *Int J Tuberc Lung Dis* 2004;8:1130–3.
 - [42] Mugusi FM, Rusizoka O, Habib N, Fawzi W. Vitamin A status of patients presenting with pulmonary tuberculosis and asymptomatic HIV-infected individuals, Dar es Salaam, Tanzania. *Int J Tuberc Lung Dis* 2003;7:804–7.
 - [43] van Lettow M, Harries AD, Kumwenda JJ, Zijlstra EE, Clark TD, Taha TE, et al. Micronutrient malnutrition and wasting in adults with pulmonary tuberculosis with and without HIV co-infection in Malawi. *BMC Infect Dis* 2004;4:6.
 - [44] Karyadi E, West CE, Schultink W, Nelwan RHH, Gross R, Amin Z, et al. A double-blind, placebo-controlled study of vitamin A and zinc supplementation in persons with tuberculosis in Indonesia: effects on clinical response and nutritional status. *Am J Clin Nutr* 2002;75:720–7.
 - [45] Pakasi TA, Karyadi E, Suratih NMD, Salean M, Darmawidjaja N, Bor H, et al. Zinc and vitamin A supplementation fails to reduce sputum conversion time in

- severely malnourished pulmonary tuberculosis patients in Indonesia. *Nutr J* 2010;9:41.
- [46] Tsukaguchi K, Okamura H, Matsuzawa K, Tamura M, Miyazaki R, Tamaki S, et al. Longitudinal assessment of IFN-gamma production in patients with pulmonary tuberculosis complicated with diabetes mellitus. *Kekkaku* 2002;77:409–13.
- [47] Walzl G, Ronacher K, Hanekom W, Scriba TJ, Zumla A. Immunological biomarkers of tuberculosis. *Nat Rev Immunol* 2011;11:343–54.
- [48] Gomez DI, Twahirwa M, Schlesinger LS, Restrepo BI. Reduced *Mycobacterium tuberculosis* association with monocytes from diabetes patients that have poor glucose control. *Tuberculosis (Edinb)* 2013;93:192–7.
- [49] Chu CM, Kuo SF, Hua CC, Wu SY, Chuang DY, Wu HP. Glucose increases interleukin-12 gene expression and production in stimulated peripheral blood mononuclear cells of type 2 diabetes patients. *Biomed J* 2014;37:293–7.
- [50] Weiner J, Parida SK, Maertzdorf J, Black GF, Repsilber D, Telaar A, et al. Biomarkers of inflammation, immunosuppression and stress with active disease are revealed by metabolomic profiling of tuberculosis patients. *PLoS One* 2012;7:e40221.

Profile of glycated-hemoglobin, antioxidant vitamin and cytokine levels in pulmonary tuberculosis patients: A cross sectional study at Pulmonary Diseases Center Semarang City, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7